

# Probabilistic procedure to estimate the macroseismic intensity attenuation in the Italian volcanic districts

G. Zonno<sup>1</sup>, R. Azzaro<sup>2</sup>, R. Rotondi<sup>3</sup>, S. D'Amico<sup>2</sup>, T. Tuvè<sup>2</sup> and G. Musacchio<sup>1</sup>



Istituto Nazionale di Geofisica e Vulcanologia  
<sup>1</sup> sezione di Milano-Pavia, Milano, Italy  
<sup>2</sup> sezione di Catania, Catania, Italy




Istituto di Matematica Applicata e Tecnologie Informatiche (CNR)  
<sup>3</sup> sezione di Milano, Milano, Italy



In Italian volcanic areas, we apply a probabilistic procedure for Macroseismic Intensity Attenuation estimates. The procedure, following the Bayesian approach, allows to exploit additional information on historical earthquakes.

The method, given the epicentral intensity and the site-epicenter distance, begins from selected earthquakes intensity data points and ends at the assessment of the intensity ( $I_s$ ) probability distribution at a site.

Our probabilistic method provides a probability function matrix that can be directly applied for the computation of probabilistic seismic hazard at the site



For the Etna region the CMT local earthquake catalogue has been used.

For the remaining Italian volcanic districts (Aeolian Islands, Vesuvius-Iscchia, and Albi Hills) the CPTI04 Italian seismic catalogue and the DBMI04 associated database have been considered.

## Mean features of the procedure

- we identify possible different decay trends through a clustering algorithm
- we estimate the probability distribution of the intensity at site  $I_s$ , conditioned on the epicenter-site distance  $d$  and on  $I_0$ , by exploiting knowledge from prior experience or data (Bayesian method)
- assuming the mode of this distribution as estimator of  $I_s$ , we forecast macroseismic fields given  $I_0$

## Step by step...

**Step 1.** Explorative analysis of a set of macroseismic fields representative of the time-space-size distribution of the Italian seismicity

**Step 2.** We divide the data set into groups/clusters of macroseismic fields that are similar to each other through an *agglomerative hierarchical clustering* based on the evaluation of the distance between each pair of rows of the matrix we have used the Manhattan distance  $d(i, j) = \sum_{k=1}^n |x_{ik} - x_{jk}|$ .

## Step 3. Probabilistic analysis of the intensity attenuation

- the intensity decay  $\Delta I$  is considered as a random variable like the macroseismic intensity.
- the variable  $\Delta I$  is discrete and belongs to the domain  $\{0, 1\}$   $\rightarrow$  it is reasonable to choose for  $I_s = I_0 - \Delta I$  at a fixed distance, the binomial distribution  $Bin(I_0, p)$  conditioned on  $I_0$  and  $p$
- ground shaking may differ even among sites located at the same distance  $\rightarrow p$  random variable


The prior hyperparameters  $\alpha, \beta$  express our initial knowledge on the decay process that we obtain by examining the three classes of macroseismic fields.

## Algorithm for estimation Given earthquakes of the same $I_0$ in class $C$

- draw  $L$  distance bins  $\{R_1, R_2, \dots, R_L\}$  around the epicenter of width  $\Delta r$
- assume that the intensity in all the sites within each  $R_j$ ,  $j = 1, \dots, L$ , band follows the same binomial distribution with parameter  $p_j$
- assign the initial value of the parameters on the basis of the macroseismic fields belonging to the same class, but with different  $I_0$
- we update the hyperparameters  $\alpha$  and  $\beta$  through its posterior mean and we estimate the parameter  $p_j$ ,  $j = 1, \dots, L$ , through its posterior mean
- in order to let the parameter of the binomial distribution for the intensity  $I_s$  at site vary with continuity, we smooth the estimates  $\hat{p}_j$ ,  $j = 1, \dots, L$ , with the method of least squares, using an inverse power function  $g(d) = (1/d)^{\gamma}$


## Step 4. Building future scenarios

- binomial distribution at any distance  $d$
- $$P_{\text{binomial}}(I_s = i | I_0 = i_0, d) = \binom{i_0}{i} g(d)^i (1 - g(d))^{(i_0 - i)}$$
- where  $g(d) = (1/d)^{\gamma}$  is the estimated inverse power function
- $\rightarrow$  the mode  $i_{\text{mode}}$  of the distribution is the predicted value



**Left Panel - Earthquakes clustering.** The 38 earthquakes of the dataset in two clusters (SET 1, SET 2). Clustering was done with the AGNES algorithm (shareware package R). The intensity dataset considered in the present analysis is the same of previous study by Azzaro et al. (2006) on the Italian volcanic districts. **Right Panel - Tab. 1: Macroseismic Intensity dataset.** Etna region: 24 events. Remaining Italian volcanic areas: Aeolian Islands (6 events), Vesuvius-Iscchia (3 events) and Albi Hills (5 events).

To summarize information in each macroseismic field, we have chosen some measures of location and dispersion of each set for epicenter-site distances with the same Macroseismic Intensity Decay: median, mean, and 3<sup>rd</sup> quartile.



## Left Panel:

Estimates of  $p_i$  (blue dots) for SET 1 and SET 2. 90% highest posterior density (HPD) intervals of the Beta distribution for each  $p_i$  (green bars). Smoothing inverse power function (red dots).

## Right Panel:

Matrix of Binomial Probability Function of intensity  $I_0 =$  VIII MCS for SET 1 and SET 2. In light blue is the mode for each bin of distance.

## Data

## Probabilistic Results


## Deterministic Results

The decay trends produced by the clustering algorithm match well the ones published in the past (Azzaro et al. 2006). This suggests that the method could be successfully applied to other cases.


## Open questions:

- Most of the earthquakes considered have epicentral intensity VII or VIII (MCS): this makes the probability functions of  $I_s$  conditioned on the other  $I_0$  (VI and IX) not very reliable.
- The method should be validated using earthquakes not included in our dataset of Table 1, on the basis of probabilistic measures of the degree to which the model predicts the decay in the data points of a macroseismic field (Rotondi and Zonno, 2004).

Probability  $\{I_s \leq i | I_0 = \text{VIII}\} > 0.50$   
 $(I_s \text{ threshold recorded with at least 50\%})$



Probability  $\{I_s > VI | I_0 = \text{VIII}\} = p$   
 $(\text{Probability of observing } I_s \text{ greater than } VI)$



**ACKNOWLEDGMENTS.** This work was partially funded by the Italian Civil Protection Department in the frame of the 2007-2009 agreement with Istituto Nazionale di Geofisica e Vulcanologia - INGV (Volcanological Project V4-Flank).

## References

- Albarelli D., Azzaro R., Barbato M.S., D'Amico V., D'Amico S., Rotondi R., Tuvè T. and Zonno G. (2007). Valutazioni di pericolosità sismica in termini di intensità macroseismica utilizzando metodi di sito. Progetto DPC-INGV S1, Deliverable D9. <http://del9.it.inpg.it/09-9.html>
- Albarelli D., D'Amico V., Gasperini P., Petrucci F., Rotondi R. and Zonno G. (2007). Nuova formulazione delle procedure per la stima dell'intensità macroseismica da dati epicentrali o da sismogrammi in zone vicine. Progetto DPC-INGV S1, <http://del9.it.inpg.it/09-9.html>
- Albarelli D. and Muccarelli M. (2002). Seismic hazard estimates using H-defined macroseismic data at site. *Pure Appl. Geophys.*, 159, 1289-1304.
- Azzaro R. and Barbato M.S. (2000). Analysis of the seismicity of Southeastern Sicily: a proposed tectonic interpretation. *Annali di Geofisica*, 43, 1, 171-188.
- Azzaro R., D'Amico S., Mostacchio A. and Scarfi L. (2006). Terremoti ed effetti macroseismici in Sicilia orientale nel periodo Gennaio 2002-Dicembre 2005. *Quad. Geofis.*, 41, pp. 62.
- Azzaro R., Barbato M.S., Rotondi R. and Rigato R. (2000). Macroseismic catalogue of Mt. Etna earthquakes from 1852 to 1998. *Atti Vulcanol.*, 12 (1), 3-36 (with CD-ROM).
- Azzaro R. and Barbato M.S. (2000). Analysis of the seismicity of Southeastern Sicily: a proposed tectonic interpretation. *Annali di Geofisica*, 43, 1, 171-188.
- Azzaro R., D'Amico S., Mostacchio A. and Scarfi L. (2002). Terremoti ed effetti macroseismici in Sicilia orientale - Calabria meridionale nel periodo Gennaio 1999 - Dicembre 2001. *Quad. Geofis.*, 47, pp. 36.

## References

- D'Amico V. and Albarelli D. (2007). Codice per il calcolo della pericolosità sismica da dati di sito: SASHA (Site Approach to Seismic Hazard Assessment). Progetto DPC-INGV S1, Deliverable D12, <http://del12.it.inpg.it/12-12.html>
- INGV-DPC 2004-2006 Agreement, Project S1 (2007). Continuation of assistance to DPC for improving and using the seismic hazard map compiled according to the Prime Minister "Ordinanza" 3274/2003 and planning future initiatives. Final Report, coord. G.M., Meletti C., Stucchi M., <http://esep1.it.inpg.it>
- Meletti C., Muccarelli M. and Albarelli D. (1994). Estimates of the seismicity rates using H-defined macroseismic data. *Rivista*, 143, 4, 617-632.
- Pasolini C., P. Gasperini, D. Albarelli, B. Loli, and V. D'Amico (2008). The Attenuation of Seismic Intensity in Italy, Part 1: Theoretical and Empirical Backgrounds. *Bull. Seism. Soc. Am.*, Vol. 98, No. 2, pp. 682-691.
- Rotondi R., Terranova A., Brambilla C. and Zonno G. (2008). The intensity attenuation of Colfiorito and other strong earthquakes: the viewpoint of forecasters and data gatherers. *Annali di Geofisica*, accepted.
- Rotondi R. and Zonno G. (2005). Seismic scenarios in terms of macroseismic intensity. *Proceedings of the First European Conference on Earthquake Engineering and Seismology*, Geneva (CH), 3-8 September 2005, Abstract, 124.
- Rotondi R. and G. Zonno (2004). Bayesian analysis of a probability distribution for local intensity attenuation. *Annali di Geofisica*, 47, 5, 1521-1540.
- Stucchi M. et al. (2007). DBMI04, il database delle osservazioni macroseismiche dei terremoti italiani utilizzate per la compilazione del catalogo parametrico CPTI04. <http://delmius.it.inpg.it/DBMI04>, Quaderno di Geofisica, 49, pp. 38.
- Zonno G., Rotondi R. and C. Brambilla (2008). Mining macroseismic fields to estimate the probability distribution of the intensity at site, submitted *Bull. Seism. Soc. Am.*
- Zonno G., Meletti C., Rotondi R. and Petrucci F. (1995). Bayesian estimation of the local intensity probability for seismic hazard assessment. *Proceeding of the Fifth International Conference on Seismic Zonation*, Nice, October 1-15, 1995, 1723-1729.